

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave Blank)	2. REPORT DATE 9 Nov 98	3. REPORT TYPE AND DATES COVERED Final 1 Oct 96 - 30 Sep 98		
4. TITLE AND SUBTITLE Turbulent Mixing in Oceanic Surface and Benthic Boundary Layers		5. FUNDING NUMBERS G N00014-97-1-0070		
6. AUTHORS H.J.S. Fernando				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Arizona Board of Regents Arizona State University, Office of Research and Creative Activities P.O. Box 871603 Tempe, AZ 85287-1603		8. PERFORMING ORGANIZATION REPORT NUMBER EFD 010		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research Ballston Tower One 800 North Quincy Street Arlington, VA 22217-5660		10. SPONSORING / MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) A comprehensive research program was carried out to investigate turbulence, waves and mixing in oceanic surface and benthic boundary layers, with special emphases on the generation, sustenance, diffusion and decay of turbulence in such layers. The research was mainly laboratory experimental, supplemented by analytical and numerical developments. The penetration of a shear-generated turbulent layer into a stably stratified non-turbulent layer, with applications to the oceanic surface-layer deepening, was the theme of the first part of the research program; the rate of turbulent-layer propagation, energetics, diffusion of mass, momentum and turbulent kinetic energy and small scale processes were the studied in detail. The applicability of results to oceanic situations, especially for surface-layer modeling, was also evaluated. The second part of the investigation dealt with oscillatory, surface-wave boundary layers, mimicked in the laboratory by an oscillatory bed underlying either a quiescent fluid layer or a mean current. The aim was to measure the diffusivities of momentum, mass and turbulent kinetic energy in such layers and to use the results to verify commonly used parameterizations in predictive models for coastal oceans.				
14. SUBJECT TERMS physical oceanography, turbulence, mixing, stratification, boundary layers, mixed layers		15. NUMBER OF PAGES 10		
		16. PRICE CODE		
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)
Prescribed by ANSI Std. Z39-1
298-102

DTIC QUALITY INSPECTED 8

**TURBULENT MIXING IN OCEANIC SURFACE AND BENTHIC BOUNDARY
LAYERS**

**Final Report of the ONR Contract No.
N00014-97-0070**

by

**H.J.S. Fernando
Department of Mechanical & Aerospace Engineering
Arizona State University
Tempe, AZ 85287-6106**

Contract Period: October 01, 1996 - September 30, 1998

Contract Monitor: Dr. Louis Goodman

EFD Report No. 010

19981125 040

Introduction

The long-term goal of our research program is to understand, using laboratory experiments, numerical modeling and theoretical analysis, small-scale mixing processes occurring in oceanic surface and benthic boundary layers. The knowledge so gained will be used to develop sound closure parameterizations for oceanic predictive numerical models. The objectives of the specific effort being reported were to improve the fundamental knowledge of turbulent mixing and diffusion processes occurring in oceanic boundary layers, with special emphases on the surface mixed layer and the wave-current boundary layer in coastal oceans. In the studies of surface mixed layers, the focus was on the penetration of a mixed layer (say, driven by the wind) into a density stratified bottom layer while transporting momentum and mass vertically. Also of interest were the effects of such transports on large-scale circulation patterns and air-sea coupling. Studies on the wave boundary layer are expected to verify the accuracy of currently used bottom boundary layer parameterizations of coastal ocean models.

During the contract period, the principal investigator H.J.S. Fernando, graduate students Rajka Krstic and Eric Strang, post-doctoral fellows Andrew Folkard and Heather Earnshaw, and visiting scientists Professors Eliezer Kit (Tel Aviv University) and J.C.R. Hunt (U.K. Meteorological Office) worked on several problems related to turbulent mixing in stably stratified fluids and coastal-ocean bottom boundary layer. The results of completed studies are outlined below and the resulting publications are listed at the end of this report.

Turbulent Mixing in Stably Stratified Shear Flows

A major part of the study on the penetration of a surface mixed-layer into an underlying stratified layer was laboratory experimental. The experiments were performed in a recirculating water channel, whereby an upper turbulent layer was driven over a stagnant denser layer to mimic the development of upper-ocean mixed layer. A uniquely designed disk pump was used to drive the flow, and special precautionary measures were used to ensure one-dimensional growth of the mixed layer. Detailed measurements using the laser-Doppler, hot film and particle-image velocimetry techniques as well as

flow imaging using the laser-induced fluorescence (LIF) method were used for flow diagnostics. The measurements included the production of turbulent kinetic energy, buoyancy flux, rate of dissipation, internal wave radiation, integral-scales of turbulence and the local Richardson number with a resolution of 2.5 mm (using a specially designed probe). The experiments were conducted by Dr. Eric Strang as a part of his Ph.D. thesis and the P.I., in collaboration with Prof. J.C.R. Hunt of the University of Cambridge, U.K, carried out a rapid-distortion-theory based analysis of the problem.

Efforts to quantify buoyancy and momentum transports through stratified shear layers using our controlled laboratory experiments revealed some interesting features. It was found that these transport rates are primarily governed by the bulk Richardson number $Ri_B = \Delta b D / \Delta U^2$, where ΔU and Δb are the velocity and buoyancy jumps across the shear layer and pycnocline, respectively, and D is the depth of the mixed layer overlying the sheared density interface. This bulk parameter, however, was closely related to the mean local gradient Richardson number $Ri_g = N^2 / (du/dz)^2$, where N is the local buoyancy frequency at the base of the mixed layer and du/dz is the local shear. When $Ri_B < 5$ (or $Ri_g < 1$), turbulent mixing at the shear layer was dominated by energetic Kelvin-Helmholtz billows, the breakdown of which produced strong, intermittent buoyancy transport episodes. Mixing in the stratified shear layer appears to be most efficient, with a mixing efficiency of $Ri_f = 0.4$, at the critical Richardson number of $Ri_g = 1$ (or $Ri_B = 5$). The range of flux Richardson numbers Ri_f observed ($0.05 < Ri_f < 0.4$) is consistent with that of $Ri_f = 0.15$ commonly employed in the Osborn dissipation model and those found in various oceanic data, namely, $Ri_f \sim 0.15 - 0.2$ in turbulent patches within the main thermocline (Moum & Osborn 1986, J. Phys. Ocean. 16, 1250) and $Ri_f \sim 0.4$ in turbulent tidal fronts (Gargett & Moum 1995, J. Geophys. Res., 81, 1180). Correspondingly, the dissipation flux coefficients (the ratio of buoyancy flux to the dissipation rate) measured in our experiments showed an excellent agreement with those obtained in turbulent tidal fronts and a reasonable agreement with data obtained from oceanic (turbulent) surface layers.

Above the critical value of $Ri_g > 1$, the mechanism responsible for turbulent mixing transitions from K-H to intermittent breaking of interfacial (internal) waves and secondary shear (Holmboe)

instabilities. Crossing to this new regime is associated with a dramatic reduction of the entrainment rate, almost by an order of magnitude. The laboratory observation of a critical Richardson number ($Ri_g = 1$) is consistent with limited ocean observations available (Moum et al. 1992, J. Phys. Ocean. 22, 1330). Furthermore, it is in fair agreement with mixing shut-off criteria employed in some integral mixed-layer models (Price et al. 1986, J. Geophys. Res., 91, 6411).

Measurements performed vertically across the stratified shear layer were used to estimate the eddy diffusivities of density and momentum, K_ρ and K_m , respectively. When properly scaled, comparisons of these values with those deduced by oceanic microstructure measurements and several expressions used for oceanic mixing parameterizations indicate a fair agreement. Laboratory data, however, were typically larger than upper bounds of these estimates. When mixing is active, $Ri_B < 5$ (or $Ri_g < 1$), K_ρ was found to be approximately equal to K_m ; this assumption is commonly used in numerical models at all Ri_B , although our data show that, at large Ri_B , the momentum transfer coefficients are higher than its buoyancy transfer counterpart.

Formation of Step Microstructure in Stratified Turbulent Flows

Numerous oceanic and atmospheric observations have provided evidence for widespread occurrence of microstructure in regions where overall stratification is stable. This microstructure can take the form of either ragged density profiles or a series of well-mixed turbulent layers separated by sheet-like density interfaces (step microstructure). No single mechanism can be attributed to all of the microstructure observations, and to this end several plausible mechanisms have been proposed. The purpose of this part of the work was to describe yet another mechanism that has not been proposed before. It was argued that turbulent mixing in a non-linearly stratified fluid or an isolated mixing event in a linearly stratified fluid can lead to the formation of step microstructure because of the merger of disparate mixing fronts. This concept is demonstrated using a simple laboratory experiment and an associated theoretical model.

Experiments on Collapsing Turbulent Regions in Stratified Fluids

Laboratory experiments were carried out to investigate the properties of a collapsing turbulent patch generated within a linearly stratified fluid by a sustained energy source and its longtime evolution in the presence of lateral boundaries. An oscillating grid spanning the width of the experimental tank was used as the turbulence source. Initially, the patch grows rapidly, as in an unstratified fluid, until the buoyancy forces arrest its vertical growth. Thereafter, the patch collapses to form horizontally propagating intrusions at its equilibrium density level. The fluid lost from the patch into the intrusion is replenished by return currents generated at the top and bottom edges of the patch. The nose of the intrusion propagates with a constant average speed ("initial spreading regime") determined mainly by the horizontal pressure gradient forces and the resistance induced by upstream propagating, low-frequency, columnar internal waves. Although the intrusion propagation speed is independent of viscous effects, they cause the development of a slug of fluid pushed ahead of the intrusion. When this slug reaches the endwall, strong upstream blocking occurs, causing the intrusion to decelerate ("blocked regime"); the intrusion nose, however, eventually reaches the endwall. The thickness of the patch is found to be approximately constant during the initial spreading regime and slowly growing in the blocked regime. At large times (t) both the patch and the "fully blocked" intrusion begin to grow vertically with a power law of the form $t^{1/5}$. A simple mixing model was advanced to explain this observation. Various turbulent and internal-wave parameters pertinent to collapsing patches were also measured, and their properties were compared with those of non-collapsing patches.

An Analysis of Turbulent Motions in and Around a Differentially Forced Density Interface

The Rapid-Distortion-Theory based analysis proposed by Fernando & Hunt [J. Fluid Mech., 347, 197, 1997] was extended to study the nature of turbulence in and around a density interface sandwiched between turbulent layers with dissimilar properties. It was shown that interfacial motions consist of low-frequency, resonantly excited, non-linear internal waves and high-frequency, linear internal waves driven by background turbulence. Based on the assumptions that (i) all resonant waves

and some non-resonant waves having frequencies close to the resonant frequencies grow rapidly, break and cause interfacial mixing, (ii) the spectral amplitude of the vertical velocity in the wave-breaking regime is constant, and (iii) kinetic energy is equipartitioned between linear and non-linear breaking wave regimes, the r.m.s. vertical velocity at the interface and the turbulent kinetic energy flux into the interface were calculated. The migration velocity of the interface was calculated using the additional assumption that the buoyancy flux into a given turbulent layer is a fixed fraction of the turbulent kinetic energy flux supplied to the interface by the same layer. The calculations were found to be in good agreement with the entrainment data obtained in previous laboratory experiments in the parameter regime where the interface is dominated by internal wave dynamics.

Studies on Turbulent Oscillatory (Wave) Boundary Layers

Two approaches were used to study the physics and transport properties of wave boundary layers. In the first approach, a purely oscillatory turbulent boundary layer, generated by an oscillating bottom in a deep fluid layer, was used to mimic the oscillatory flow under waves. This work was performed by Ms. Rajka Krstic as a part of her Ph.D. thesis project. In the second study, carried out by Dr. Heather Earnshaw, an actual wave boundary layer produced by a periodic train of waves traveling on a sloping beach was investigated in a large wave tank of dimensions 104.5 x 3.5 x 6 (ft). In both studies, the parameters of interest are the mean velocity profiles, integral length and velocity scales of turbulence and the upward diffusion of turbulent kinetic energy, momentum and mass. Particle-image, particle-tracking and laser-Doppler velocimetry are used for flow diagnostics.

Flow visualization studies during the oscillatory boundary layer work elicited the mechanism of vortex formation surrounding roughness elements, which occurred at the end of each half-cycle. Vortices so formed interacted with surrounding vortices within the boundary layer, causing violent exchange of the mass and momentum in the vertical direction. Turbulent kinetic energy was found to be maximum near the bed, and its distribution followed the numerical calculations of Justensen (1988, Coastal Engineering, 12, 257) reasonably well. Measurements over two-dimensional planes enabled the calculation of autocorrelation functions of velocity at

different distances from the bed for different phases of flow oscillations. Comparison of eddy viscosity measurements with the theoretical model of Trowbridge and Madsen (1984, J. Geophys. Res., 89, 7989) and the integral length scale measurements with the model of Grant & Madsen (1979, J. Geophys. Res., 80, 5109) elicited successes and failures of these models. For example, integral length-scale measurements agreed fairly well with the predictions of Grant & Madsen, usually within 10%, indicating their suitability to parameterize lengthscales in numerical models. As a continuation of this work, experiments are now being conducted on wave-current boundary layers. A specially designed apparatus, capable of superimposing a steady (mean) current on the oscillatory motion is being used for this purpose. This study is expected to yield important new information on the modification of current (mean velocity, shear stresses, roughness length and integral velocity and lengthscales) by the oscillatory component of the boundary layer. The results will be compared with the theoretical model of Grant & Madsen.

List of Publications

Journal Papers

- DeSilva, I.P.D., Fernando, H.J.S., Eaton, F. and Hebert, D., "Kelvin-Helmholtz Billows in Nature and laboratory," *Earth & Planetary Sciences Letters*, **143** (1-4), 217-231, 1996.
- Kit, E., Strang, E. and Fernando, H.J.S., "Measurement of Turbulence Near Shear-Free Density Interfaces," *Journal of Fluid Mechanics*, **334**, 293-314, 1997.
- Voropayev, S.A., Zhang, X., Boyer, D.L., Fernando, H.J.S. and Wu, A., "Horizontal Jets in Rotating Stratified Fluids," *Physics of Fluids*, **9**(1), 115-126, 1997.
- Kit, E. and Fernando, H.J.S., "Frequency Spectra of Scalar Fluctuations at Entraining Stratified Interfaces," *Fluid Dynamics Research*, **19**, 65-75, 1997.
- Folkard, A., Davies, P.A. and Fernando, H.J.S., "Measurements of Turbulent Patch in a Rotating Linearly Stratified Fluid," *Dynamics of Atmospheres and Oceans*, **26**, 27-51, 1997.
- Fernando, H.J.S. and Hunt, J.C.R., "Turbulent Mixing Across Shear-Free Density Interfaces; Part 1 - Modeling Considerations," *Journal of Fluid Mechanics*, **347**, 197-234, 1997.
- McGrath, J., Fernando, H.J.S. and Hunt, J.C.R., "Turbulent Mixing Across Shear-Free Density Interfaces; Part 2 - Laboratory Experiments," *Journal of Fluid Mechanics*, **347**, 235-261, 1997.
- Zhang, X., Boyer, D.L. and Fernando, H.J.S., "Turbulence-Induced Rectified Flows in Rotating Fluids," *Journal of Fluid Mechanics*, **350**, 97-118, 1997.

- DeSilva, I.P.D. and Fernando, H.J.S., "Experiments on Collapsing Patches in Stratified Fluids," *Journal of Fluid Mechanics*, **358**, 29-60, 1998.
- Manasseh, R., Ching, C.Y. and Fernando, H.J.S. "The Transition from Wave-Dominated to Density Driven Isolated Flows," *Journal of Fluid Mechanics*, **361**, 253-274, 1998.
- Luccio, P.A., Voropayev, S.A., Fernando, H.J.S., Boyer, D.L. and Houston, W.N. "The Motion of Cobbles in the Swash Zone," *Coastal Engineering*, **33**, 41-60, 1998.
- Fernando, H.J.S., Chen, R-r. and Ayotte, B.A., "Development of a Point Plume in the Presence of Background Rotation," *Physics of Fluids*, **10(9)**, 2369-2383, 1998.
- Voropayev, S., Roney, J., Boyer, D.L., Fernando, H.J.S. and Houston, W.N. "The Motion of Large Bottom Particles (Cobbles) in Wave-induced Oscillatory Flows," *Coastal Engineering*, **34**, 197-219, 1998..
- Fonseka, S.V., Fernando, H.J.S. and van Heijst, G.J.F., "The Evolution of an Isolated Turbulent Region in a Stratified Fluid," *Journal of Geophysical Research* (accepted for publication).
- Colomer, J. Casamitjana, X. and Fernando, H.J.S. "Resuspension and Sedimentation of Particles from a Sediment Bed by Turbulent Jets," *Applied Scientific Research*, (accepted for publication).
- Srdic-Mitrovic, A.N., Mohammed, N.A, and Fernando, H.J.S., "Gravitational Settling of Particles Through Density Interfaces," *Journal of Fluid Mechanics* (accepted for publication).

Papers Submitted

- Strang, E.J. and Fernando, H.J.S. "Turbulence and Mixing in Stratified Shear Flows," *Journal of Fluid Mechanics* (under revision).
- Neves, J. and Fernando, H.J.S., "Behavior of a Sediment-Laden Vertical Turbulent Jet," *Journal of Fluid Mechanics* (under revision).
- Lozovatsky, I. and Fernando, H.J.S., "Observations of Turbulence in Marine Stratified Layers," *Journal of Geophysical Research* (under revision).
- Wu, A. and Fernando, H.J.S, "An Analysis of Fluid Motions in and Around Differentially Forced Density Interface," *Theoretical & Computational Fluid Mechanics* (under revision).
- Fernando, H.J.S. and Buhler, J., "The Scales of Sediment Fingers," *Journal of Marine Research* (under revision).
- Colomer, J. and Fernando, H.J.S. Resuspension of Sediments by Multiple Vertical Jets," *ASCE Journal of Hydraulic Engineering* (under revision).
- DeSilva, I.P.D., Brandt, A., Montenegro, L.M. and Fernando, H.J.S. Gradient Richardson Number Measurements in a Stratified Shear Layer. Submitted for Publication.
- Voropayev, S.A., McEachern, B., Fernando, H.J.S. and Boyer, D.L. "Large Vortex Structures Behind Maneuvering Bodies in Stratified Fluids," Submitted for Publication.

Conference Proceedings

- Fernando, H.J.S., Colomer, J. and Boubnov, B. "Laboratory Studies of Convection from Isolated Sources," *Proceedings, Sixth Asian Congress of Fluid Mechanics*, Madras, 857-859, 1997.
- Strang, E., Fernando, H.J.S. and Kit, E. "Aspects of Turbulent Mixing at Density Interfaces," *Proceedings, Fifth IMA Conference on Stratified Flows - "Mixing and Dispersion in Stably Stratified Flows,"* Accepted, Oxford University Press, 1998.
- Hunt, J.C.R. and Fernando, H.J.S. "Separated Flows Around Bluff Obstacles at Low Froude Numbers; Vortex Shedding and Estimates of Drag," *Proceedings, Fifth IMA Conference on Stratified Flows - "Mixing and Dispersion in Stably Stratified Flows,"* Accepted, Oxford University Press, 1998.
- Folkard, A.M. and Fernando, A., "How is Thorpe-Scale Decay Rate Related to External Parameters? Physical and Theoretical Studies," *Proceedings, Fifth IMA Conference on Stratified Flows - "Mixing and Dispersion in Stably Stratified Flows,"* Accepted, 1998.
- Kit, E., Strang, E., Fernando, H.J.S. and Norton, H. "Experiments on the Development of K-H billows in stratified shear flows," *Seventh European Turbulence Conference, In: Advances in Turbulence VII* (Ed. Uriel Frisch), Kluwer Academic Publishers, 1998.
- Mahalov, A., Nicolaenko, B. and Fernando, H.J.S. "Development and Verification of Experimental Data Bases for Pancake Structures in Atmospheric Flows," *Proceedings 12th International Symposium on Aerospace/Defense Sensing, Simulation and Controls*, Apr 13-17, Orlando, 1998.
- Srdic-Mitrovic and Fernando, H.J.S. "On the Stratification Drag," *Second International Symposium on Environmental Hydraulics*, Hong-Kong, Dec 16-18, Accepted, 1998.
- Lozovatsky, I.D. and Fernando, H.J.S. "Turbulence in Stratified Patches on a Shallow Shelf," *Proceedings of the Konstantin Federov Memorial Symposium*, St. Petersburg., UNESCO Volume # x, in Press.

Conference Presentations

- Fernando, H.J.S. "Modeling of Geophysical Flows," Invited Speaker, Moscow'97 Conference on Stability and Instabilities of Stratified and Rotating Flows," Moscow, 1997.
- Fernando, H.J.S. "Laboratory Experiments on Stratified Shear Flows," Keynote Speaker, International Workshop on Mixing in Geophysical Flows, Barcelona, Spain, 1997.
- Fernando, H.J.S. and Corder, D. "The Evolution of a Turbulent Patch in a Stratified Shear Flow," *Bull. Am. Phys. Soc.* **42(11)**, 2200, 1997.
- Srdic-Mitrovic, A., and Fernando, H.J.S. "The Influence of Background Turbulence on the Settling Velocity of Heavy Particles," *Bull. Am. Phys. Soc.* **42(11)**, 2216, 1997.
- Casamitjana, X., Colomer, J., and Fernando, H.J.S. "Resuspension of a Particle Bed in a Cone-Shaped Basin." *Applications to Lake Banyoles, XXXVII SIL Congress*, 9-14 August, University College of Dublin, 1998.
- Strang, E.J. and Fernando, H.J.S. "Shear Induced Vertical Mixing and Transport in the Upper Ocean," *Am. Geophys. Union, Ocean Sci., EOS*, **79(1)**, 152, 1998

- Norton, H., Strang, E., Kit, E. and Fernando, H.J.S. "Distortion of Turbulent Regions by Stratified Shear Flows," *Am. Geophys. Union, Ocean Sci., EOS*, **79(1)**, 152, 1998
- Earnshaw, H.C. and Freehand, H.J.S. " Oscillatory Bottom Boundary Layers Induced on a Shallow Break," *Am. Geophys. Union, Ocean Sci., EOS*, **79(1)**, 152, 1998
- McEachern, E.B., Voropayev, S.I., Boyer, D.L. and Fernando, H.J.S. "Laboratory Modeling of Wakes Behind Maneuvering Self-Propelled Bodies and Related Vortex Structures ion Stratified Fluids," *Am. Geophys. Union, Ocean Sci., EOS*, **79(1)**, 152, 1998
- Fernando, H.J.S. "Turbulence and Mixing in Environmental Flows," Invited Paper, "Hopkins Conference in Environmental Fluid Mechanics," 8-9, April 2-4, Baltimore, MD 21218, 1998.
- Fernando, H.J.S. "Stratified Turbulence," Invited Pedagogical Lecture, IUTAM, IAMAS/IUGG, GTP/NCAR Workshop on "Developments in Geophysical Turbulence," Boulder, CO June 16-19, 1998.
- Fernando, H.J.S. Entrainment and Mixing in Stably Stratified Flows, Invited Lecture, GTP/NCAR conference on "Observations, Experiments and LES" Boulder, CO Aug. 12-15, 1998.
- Mahalov, A., Berman, N.S., Fernando, H.J.S., Yu, F. and Pardyjak, E. "Stable Layers in the Atmospheric Boundary Layer," *Bull. Am. Phys. Soc.* **xx(11)**, xxxx, 1998.